



## Role of Free Radicals and Antioxidants in Mastitis

Mahmoud R. Abd Ellah\*

*Department of Animal Medicine, Faculty of Veterinary Medicine, Assiut University 71526, Egypt*

Mastitis is a worldwide problem and of major economic threat to dairy farmers. Its incidence depends on the microorganisms, the surrounding environment and the defense mechanisms in the udder tissues and blood. In the dairy industry, both clinical and subclinical mastitis produce great economic losses. Subclinical mastitis has no apparent clinical signs but accompanied with elevation of somatic cells count (SCC) in milk. However, clinical mastitis has elevated body temperature, inappetance, a red, swollen and/or painful udder and/or abnormal milk (Huijps *et al.*, 2008). Milk with higher SCC is positively associated with malondialdehyde level in milk (Atroshi *et al.*, 1996; Cesa, 2004; Miranda *et al.*, 2004; Suriyasathaporn *et al.*, 2006), consequently more free radicals being released and a state of oxidative stress arise (Su *et al.*, 2002). The inflammatory reaction accompanying mastitis is generally caused by bacterial infection and results in damage to the secretory epithelium of the mammary gland and the release of cytotoxic radicals and pro-inflammatory cytokines by the phagocytic cells (Knaapen *et al.*, 1999).

### Oxidative stress in mastitis

Oxidative stress is associated with some disorders of dairy cattle as mastitis (Erskine *et al.*, 1987; Parantainen *et al.*, 1987), retained placenta and udder edema (Miller *et al.*, 1993). Mastitis could induce increase formation of free radicals in milk leading to oxidative stress (Gu *et al.*, 2009) especially during the early lactation period (Bernabucci *et al.*, 2005; Castillo *et al.*, 2005; Sordillo *et al.*, 2007). Both clinical and subclinical mastitis are associated with release of free radicals, increased total oxidant capacity and decreased total antioxidants capacity in milk (Atakisi *et al.*, 2010).

There are some important factors that contribute to the voluminous production of free radicals in mastitic milk; one of these factors is the increased numbers of cells like neutrophils, macrophages, lymphocytes, eosinophiles and various epithelial cells of mammary tissue (Smith, 1994; Knaapen *et al.*, 1999). Another factor is the increase of some cytokines (TNF- $\alpha$ , IL-1b, IL-6, IL-8) and other molecules such as nitric oxide (NO $\cdot$ ) (Blum *et al.*, 2000; Riollet *et al.*, 2000; Notebaert *et al.*, 2008). Cytokines activate reactive nitrogen intermediates (RNI), which play a complex role in the inflammatory process (Moilanen and Vapaatalo, 1995; Goff *et al.*, 1996; MacMicking *et al.*, 1997).

One of the most important RNI is NO $\cdot$ , which operates in a variety of tissues to regulate a diverse range of physiological processes, including the inflammatory response (Dawson and Dawson, 1995). During inflammation, macrophage and epithelial cells of the mammary gland produce a significant amount of NO $\cdot$ ; this inducible NO $\cdot$  mediates inflammation during mastitis (Bouchard *et al.*, 1999). Another source for NO $\cdot$  is the mammary epithelial cells and/or mononuclear phagocytes, which contribute to NO $\cdot$  production upon stimulation with lipopolysaccharide and cytokines (Boulanger *et al.*, 2001).

Nitric oxide has a key role in mediating microbistatic and/or microbicidal activity, as the activated macrophages synthesize NO $\cdot$  (Jungi, 2000), which is considered as a primary defence system that eliminate intracellular pathogens (Huie and Padmaja, 1993; O'Flaherty *et al.*, 2003). The antimicrobial properties of NO $\cdot$  is attributed to peroxynitrite, a reactive nitrogen metabolite, derived from oxidation of NO $\cdot$  (Beckman *et al.*, 1990). In severe mastitis, peroxynitrite is produced in excess, which may result in alterations in the antioxidant balance (Chaiyotwittayakun *et al.*, 2002). This means that excessive release of NO $\cdot$  results in oxidative damage to mammary gland secretions

\*Corresponding author: M.R. Abd Ellah

E-mail address: mrushdi@aun.edu.eg

(Bouchard *et al.*, 1999; Komine *et al.*, 2004; Atakisi *et al.*, 2010).

Nitric oxide was reported to increase in milk during clinical mastitis (Lacasse *et al.*, 1997, Atakisi *et al.*, 2010) and in milk and plasma after intramammary infusion of *E. coli* bacteria or *E. coli* endotoxin (Bouchard *et al.*, 1999; Blum *et al.*, 2000; Komine *et al.*, 2004). Milk SCC is a well-known indicator reflecting mammary health and milk quality, a positive correlation between SCC and NO $\cdot$  concentration had been reported (Bouchard *et al.*, 1999; Atakisi *et al.*, 2010). Changes in milk NO $\cdot$  level thus could be proposed as an additional diagnostic tool to detect inflammation during mastitis. It is clear from the above paragraphs that the body uses free radicals to destruct microorganisms at the site of inflammation. However, with chronicity and severity of inflammation more free radicals being released at the site of inflammation, which accompanied with oxidative damage to tissues, lipid peroxidation and oxidative stress.

### Protective role of antioxidants in mastitis

The cells contain a variety of antioxidants that play an important role in the protection against excessive release of reactive oxygen species in blood and tissues (Pár and Jávora, 1984; Halliwell, 1991) including the udder tissue and milk in mastitis (Smith *et al.*, 1984; Erskine, 1993; Bowers, 1997; Weiss *et al.*, 1997; Scaletti *et al.*, 2003). Antioxidants protect the body from free radicals either by directly scavenging free radicals or by inhibiting the activity of oxidizing enzymes (Abd Ellah *et al.*, 2009; Abd Ellah, 2010). The antioxidant system consists of antioxidant enzymes (superoxide dismutase (SOD), catalase, and glutathione peroxidase (GSH-Px), glutathione, ancillary enzymes (glutathione reductase (GR), glutathione S-transferase, and glucose 6-phosphate dehydrogenase, metal-binding proteins (transferrin, ceruloplasmin, and albumin), vitamins (alpha-tocopherol, ascorbate, and beta-carotene), flavonoids, and urate (Halliwell, 1994; Abd Ellah, 2010).

It had been reported that antioxidant supplementation could decrease the duration, incidence, and severity of clinical mastitis (Smith *et al.*, 1984). Despite significant evidence supporting the role of antioxidants in enhancing resistance to mastitis,

there is no information as to how the expression of specific antioxidant defenses change in mammary tissues during the preparturient period when dairy cattle experience increased oxidative stress.

In dairy cattle, cytosolic GSH-Px is the predominant intracellular form of GSH-Px and it is the most studied selenoprotein (Smith *et al.*, 1997). The activity of GSH-Px in milk varies with the species and diet (Fox and Kelly, 2006). Glutathione peroxidase catalyzes the reduction of various peroxides, protecting the cell against oxidative damage (Abd Ellah *et al.*, 2007) and protects milk lipids from oxidation (Bhattacharya *et al.*, 1988). GSH-Px activity is often used as a diagnostic tool when assessing the selenium status of dairy cows or as an indicator of increased ROS accumulation (Gärtner, 2009; Köhrle and Gärtner, 2009). It has been reported that blood concentration of selenium and GSH-Px activity negatively correlated with the prevalence of intramammary infection (Erskine *et al.*, 1987). Low GSH-Px activity in mastitic cows with high SCCs and high level of prostaglandin formation has been recorded (Hogan *et al.*, 1993), which may be attributed to the excessive release of free radicals that may result in inhibiting enzymes activity.

Superoxide dismutase was reported first in milk by Hicks *et al.* (1975), and they suggested that SOD may play an important role in keeping the oxidative stability of milk (Holbrook and Hicks, 1978). The activity of milk SOD varies between breeds of cows (Holbrook and Hicks, 1978; Lindmark-Mansson and Akesson, 2000). In cow's milk, it is not affected by stage of lactation or age of cow, and/or high SCC (Przybylska *et al.*, 2007). Based on previous studies, Atroshi *et al.* (1986) reported a substantial decrease in blood SOD in cows with mastitis, while Cetin *et al.* (2005) reported an increase in erythrocyte SOD activity in ewes with gangrenous mastitis, which may be attributed to the elevated need of the enzyme to boost the defensive mechanism of the animal against oxidation.

During inflammatory diseases, high levels of NO $\cdot$  react with superoxide anions leading to formation of peroxynitrite radical (Beckman *et al.*, 1990), the latter oxidize long chain fatty acids in cell membranes leading to lipid peroxidation (Al-Sa'doni and Ferro, 2000; Wang *et al.*, 2002), which may inhibit the activity of some antioxidant molecules as GSH-Px (Atroshi *et al.*, 1996) leading to oxidative stress (Goff *et al.*, 1996; Komine *et al.*,

2004; Weiss *et al.*, 2004).

## Role of micronutrients antioxidants in mastitis

Micronutrients that are associated with antioxidant activity including vitamin A, vitamin E,  $\beta$ -carotene, selenium, zinc, and copper had been studied for their effect on mastitis (Erskine, 1993; Bowers, 1997; Mukherjee, 2008). Supplementation of antioxidant vitamins may reduce oxidative stress and the inflammatory response during mastitis. In dairy cows, supplementation with vitamin E (Smith *et al.*, 1984) or a combination of vitamins A, D, and E (Barnouin and Chassagne, 1998) played a significant role in reducing the incidence of clinical mastitis.

Ascorbic acid is the most important water-soluble antioxidant in mammals (Saubertlich, 1994). It has been identified as oxidative stress biomarkers in bovine mastitis. Many studies had been reported that its milk concentration significantly decreased in acute and subclinical forms of mastitis especially when accompanied by an increase in the levels of lipid hydroperoxide in erythrocytes (Weiss *et al.*, 2004; Kleczkowski *et al.*, 2005, Ranjan *et al.*, 2005).

Vitamin C administered to cows by subcutaneous injection may have therapeutic value in mastitis (Chaiyotwittayakun *et al.*, 2002; Naresh *et al.*, 2002; Ranjan *et al.*, 2005). However, its therapeutic effect decreased in the presence of lipid peroxidation with moderate improvement in clinical signs of mastitis (Ranjan *et al.*, 2005).

Vitamin E and selenium (Se) are fundamental components of the antioxidant defence of tissues and cells. Vitamin E is the most important lipid-soluble antioxidant and the biologically most active form is  $\alpha$ -tocopherol. Vitamin E is an integral component of all lipid membranes and has a role in protecting lipid membranes from attack by high tissue concentrations of reactive oxygen species (Rice and Kennedy, 1988).

Vitamin E deficiencies are frequently observed in peripartum dairy cows. Most cases of clinical mastitis occur during the first month of lactation (Green *et al.*, 2002), and coincide with the lowest vitamin E blood concentration (Goff and Stabel, 1990). It was suggested that maintaining an optimal vitamin E level, together with low levels of oxidative stress is an important factor in dry cow man-

agement and improvement of udder health.

Dairy cattle have several known endogenous antioxidant defense mechanisms that can counteract the harmful effects of ROS accumulation. The selenium-dependent selenoproteins had been studied extensively with respect to mammary gland health (Miller *et al.*, 1993). Selenium supplementation to periparturient cows reduces the incidence and severity of mastitis (Smith *et al.*, 1984). The beneficial effects of selenium supplementation are thought to be due to the actions of certain antioxidant selenium-dependent enzymes (Papp *et al.*, 2007). Selenium is an integral component of the enzyme GSH-Px (Diplock, 1981; Erskine, 1993; Smith *et al.*, 1997). Several recent studies, documented the presence of other selenoprotein enzymes in the blood and tissues of dairy cattle that may play an important role in controlling oxidative stress, including thioredoxin reductase and phospholipid hydroperoxide glutathione peroxidase (Trigona *et al.*, 2006; Balla *et al.*, 2007; Bruzelius *et al.*, 2007; Sordillo *et al.*, 2007).

Pavalata *et al.* (2002) found that about 50% of dairy cows sampled in the Czech Republic had marginal or deficient Se status. Other studies in Slovenia, Germany and Ireland have reported similar findings with the majority of dairy cows being marginal or deficient in Se (Mee *et al.*, 1993; Gründer and Auer, 1995; Züst *et al.*, 1996). Weiss *et al.* (1990) found that high serum Se concentrations were associated with reduced rates of clinical mastitis and low milk SCC. Erskine *et al.* (1987) found that herds with the low SCC (less than or equal to 150,000 cells/ml) had significantly higher mean blood Se-dependent GSH-Px activity than did the herds with the high SCC (greater than or equal to 700,000 cells/ml). Whole blood concentrations of Se were also significantly higher in the herds with low SCC than in the herds with high SCC.

Deficiencies in either vitamin E or Se have been related to increased incidence and severity of mastitis. Reduced neutrophils activity is a known physiological consequence of  $\alpha$ -tocopherol or selenium deficiency. Vitamin E and the selenium-containing enzyme protect neutrophils from the destructive action of toxic oxygen molecules, which is necessary for intracellular kill of ingested pathogens. Recently, trials have shown that subcutaneous injections of vitamin E approximately 10 days prior to calving successfully raised neutrophil's  $\alpha$ -tocopherol concentrations during

the preparturient period and negated the suppressed intracellular kill of bacteria by neutrophils that are commonly observed at calving (Hogan *et al.*, 1990; Hogan *et al.*, 1993). Furthermore, dietary supplementation of cows with Se and Vitamin E resulted in a more rapid PMN influx into milk following intramammary bacterial challenge and increased intracellular killing of ingested bacteria (Smith *et al.*, 1997), as well as lowering the frequency and shortening the duration of clinical mastitis (Smith *et al.*, 1984).

Substantial research has been carried out on the effect of selenium and vitamin E on the immune function of the mammary gland and subsequent improvement in bovine udder health and mastitis control (Cerri *et al.*, 2009; Salman *et al.*, 2009). Vitamin E and selenium given prior to calving have been proposed for prophylactic treatment of mastitis (Smith *et al.*, 1997). Cows supplemented with vitamin E throughout the dry period had lower incidence of clinical mastitis during the next lactation. However, cows supplemented with both vitamin E and Se had shorter duration of clinical signs than cows supplemented with either micronutrient alone (Smith *et al.*, 1984). In a study by Smith *et al.* (1985), diets of heifers supplemented with Se and vitamin E from 60 days pre-partum and continuing throughout lactation had significantly fewer infected quarters at calving, reduced prevalence of infection throughout lactation, fewer cases of clinical mastitis, infections of shorter duration, and lower milk SCC compared with the none supplemented heifers.

Both  $\beta$ -carotene and vitamin A play an important role in protecting udder tissue and milk from the harmful effect of free radicals. Beta-carotene functions independently of vitamin A in mastitis and reproduction (McDowell, 1989). It was reported that, low concentrations of plasma vitamin A (<80 $\mu$ g/100 ml) and  $\beta$ -carotene (<200 $\mu$ g/100 ml) were linked with severity of mastitis (Chew *et al.*, 1982; Jukola *et al.*, 1996). Dahlquist and Chew (1985) carried out a study, the results revealed that cows fed vitamin A plus  $\beta$ -carotene had fewer new infections during the early dry period than cows fed the other treatments. Chew and Johnston (1985) carried out another study and found that vitamin A plus  $\beta$ -carotene reduced mean SCC during the 2nd to the 8th weeks of lactation. In another study, supplementation of vitamin A and/or  $\beta$ -carotene for cows did not affect new infections during the dry

period, new infections at calving and cases of clinical mastitis (Oldham *et al.*, 1991).

Minerals (Zinc, copper, iron, manganese, selenium) are essential for the formation of antioxidants' enzymes. For example, zinc is required for the formation of Mn-Zn SOD, deficiency of zinc affect the activity of SOD in blood and tissues, which results in increased superoxide radicals. It is known that mastitis associated with increased SCC counts in milk, which act as a source for free radicals and hence oxidative stress. Supplementation of zinc helps the animal to recover from increased oxidative stress by reducing the SCC (Kincaid *et al.*, 1984; Popovic, 2004)). Experimental studies approved that copper supplementation reduced the severity of clinical signs of *E. coli* mastitis (Scaletti *et al.*, 2003). In a separate study, Scaletti *et al.* (2002) attempted to determine the effect of copper source on the response to intra-mammary challenge with *E. coli* at day 32 of lactation. The study revealed that copper and zinc supplementation to mastitic cows enhanced the resistance to bacterial mastitis.

## Conclusion

The fact that mastitis paralyze the milk production of dairy animals, make that disease of major importance. Another fact is that the increased free radicals during mastitis are mostly associated with decreased antioxidants defence as established by many researchers. This makes the supplementation of mastitic animals with nutrients antioxidants as vitamin C, E and beta carotene, and minerals very important to help the animal recover early. On the other hand, diary animals must be supplemented with the optimal ration that contains all required nutrients needed for optimal production and reproduction. Antioxidants status of milk could be used as a diagnostic indicator for early detection of mastitis. Another possible use for measuring antioxidants in milk is the evaluation of treatment and also establishing a prognosis.

## References

- Abd Ellah, M.R., 2010. Involvement of free radicals in animal diseases. *Comparative Clinical Pathology* 19, 615–619.
- Abd Ellah, M.R., Okada, K., Goryo, M., Oishi, A., Yasuda, J., 2009. Superoxide dismutase activity as a measure of hepatic oxidative stress in cattle following ethionine

- administration. The Veterinary Journal 182, 336-341.
- Abd Ellah, M.R., Yasuda, J., Okada, K., 2007. Oxidative stress and bovine liver diseases: Role of glutathione peroxidase and glucose 6-phosphate dehydrogenase. Japanese Journal of Veterinary Research 54, 163-173.
- Al-Sa'doni, H., Ferro, A., 2000. S-nitrosothiols, a class of nitric oxide-donor drugs. Clinical Science 98, 507-520.
- Atakisi, O., Oral, H., Atakisi, E., Merhan, O., Metin Pancarci, S., Ozcan, A., Marasli, S., Polat, B., Colak, A., Kaya, S., 2010. Subclinical mastitis causes alterations in nitric oxide, total oxidant and antioxidant capacity in cow milk. Research in Veterinary Science 89, 10-13.
- Atroshi, F., Parantainen, J., Sankari, S., Jarvinen, M., Lindberg, L.A., Saloniemi, H., 1996. Changes in inflammation-related blood constituents of mastitic cows. Veterinary Research 27, 125-135.
- Atroshi, F., Parantainen, J., Sankari, S., Osterman, T., 1986. Prostaglandin and glutathione peroxidase in bovine mastitis. Research in Veterinary Science 40, 361-366.
- Balla J., Vercellotti G.M., Jeney V., Yachie A., Varga Z., Jacob H.S., 2007. Heme, heme oxygenase, and ferritin, How the vascular endothelium survives, and dies. in an iron-rich environment. Antioxidants Redox Signal 9, 2119-2137.
- Barnouin, J., Chassagne, M., 1998. Factors associated with clinical mastitis incidence in French dairy herds during late gestation and early lactation. Veterinary Research 2, 159-171.
- Beckman, J.S., Beckman, T.W., Chen, J., Marshall, P.A., Freeman, B.A., 1990. Apparent hydroxyl radical production by peroxynitrite, implications for endothelial injury from nitric oxide and superoxide. Proceedings of the National Academy of Sciences of the United States of America 87, 1620-1624.
- Bernabucci, U., Ronchi, B., Lacetera, N., Nardone, A., 2005. Influence of body condition score on relationships between metabolic status and oxidative stress in periparturient dairy cows. Journal of Dairy Science 88, 2017-2026.
- Bhattacharya, I.D., Picciano, M.F., Milner, J.A., 1988. Characteristics of Human Milk Glutathione Peroxidase. Biological Trace Element Research 18, 59-70.
- Blum, J.W., Dosogne, H., Hoeben, D., Vangroenweghe, F., Hammon, H.M., Bruckmaier, R.M., Burvenich, C., 2000. Tumor necrosis factor-(alpha) and nitrite/nitrate responses during acute mastitis induced by Escherichia coli infection and endotoxin in dairy cows. Domestic Animal Endocrinology 19, 223-235.
- Bouchard, L., Blais, S., Desrosiers, C., Zhao, X., Lacasse, P., 1999. Nitric oxide production during endotoxin-induced mastitis in the cow. Journal of Dairy Science 82, 2574-2581.
- Boulanger, V., Bouchard, L., Zhao, X., Lacasse, P., 2001. Induction of Nitric oxide production by bovine mammary epithelial cells and blood leukocytes. Journal of Dairy Science 84, 1430-1437.
- Bowers, T.L., 1997. Nutrition and immunity part 2, the role of selected micronutrients and clinical significance. Veterinary Clinical Nutrition 4, 96-101.
- Bruzelius, K., Hoac, T., Sundler, R., Onning, G., Akesson, B., 2007. Occurrence of selenoprotein enzyme activities and mRNA in bovine mammary tissue. Journal of Dairy Science 90, 918-927.
- Castillo, C., Hernandez, J., Bravo, A., Lopez-Alonso, M., Pereira, V., Benedito, J.L., 2005. Oxidative status during late pregnancy and early lactation in dairy cows. Veterinary Journal 169, 286-292.
- Cerri, R.L., Rutigliano, H.M., Lima, F.S., Araújo, D.B., Santos, J.E., 2009. Effect of source of supplemental selenium on uterine health and embryo quality in high-producing dairy cows. Theriogenology 71, 1127-1137.
- Cesa, S.J., 2004. Malondialdehyde contents in infant milk formulas. Journal of Agricultural and Food Chemistry 52, 2119-2122.
- Cetin, H., Gurgoze, S.Y., Keskin, O., Atli, M.O., Korkmaz, O., 2005. Investigation of antioxidant enzymes and some biochemical parameters in ewes with gangrenous mastitis. Turkish Journal of Veterinary and Animal Sciences 29, 303-308.
- Chaiyotwittayakun, A., Erskine, R.J., Bartlett, P.C., Herdt, T.H., Sears, P.M., Harmon, R.J., 2002. The effect of ascorbic acid and L-histidine therapy on acute mammary inflammation in dairy cattle. Journal of Dairy Science 85, 60-67.
- Chew, B.P., Hollen, L.L., Hillers, J.K., 1982. Relationship between vitamin A and beta-carotene in blood plasma and milk and mastitis in Holsteins. Journal of Dairy Science 65, 2111-2118.
- Chew, B.P., Johnston, L.A., 1985. Effects of supplemental vitamin A and beta-carotene on mastitis in dairy cows. Journal of Dairy Science 68 (supp. 1), 191.
- Dahlquist, S.P., Chew, B.P., 1985. Effects of vitamin A and beta-carotene on mastitis in dairy cows during the early dry period. Journal of Dairy Science 68 (supp. 1), 191.
- Dawson, T.M., Dawson, V.L., 1995. Nitric oxide, actions and pathological roles. Neuroscientist 1, 7-18.
- Diplock, A.T., 1981. The role of vitamin E and selenium in the prevention of oxygen-induced tissue damage. In, Selenium in Biology and Medicine. p 303. Spallholz JE, Martin JL and Ganther HE, eds.. AVI Publishing, Hartford, CT.
- Erskine, R.J., 1993. Nutrition and mastitis. Update on bovine mastitis. Veterinary Clinics of North America Food Animal Practice 9, 551-556.
- Erskine, R.J., Eberhar, R.J., Hutchinson, L.J., 1987. Blood selenium concentrations and glutathione peroxidase activities in dairy herds with high and low somatic cell counts. Journal of the American Veterinary Medical Association 11, 1417-1421.
- Fox, P.F., Kelly, A.L., 2006. Indigenous enzymes in milk, Overview and historical aspects-Part 2. International Dairy Journal 16, 517-532.
- Gärtner, R., 2009. Selenium and thyroid hormone axis in critical ill states, an overview of conflicting view points. Journal of Trace Elements in Medicine and Biology 2, 71-74.
- Goff, J.P., Stabel, J.R., 1990. Decreased plasma retinol, alpha-tocopherol, and zinc concentration during the periparturient period, Effect of milk fever. Journal of Dairy Science 73, 3195-3199.
- Goff, W.L., Carl, Johnson, W., Wyatt, C.R., Cluff, C.W.,

1996. Assessment of bovine mononuclear phagocytes and neutrophils for induced L-arginine-dependent nitric oxide production. *Veterinary Immunology and Immunopathology* 55, 45–62.
- Green, M.J., Green, L.E., Medley, G.F., Schukken, Y.H., Bradley, A.J., 2002. Influence of dry period bacterial intramammary infection on clinical mastitis in dairy cows. *Journal of Dairy Science* 85, 2589-2599.
- Gründer, H.D., Auer, S., 1995. Selenium supply of cattle in Hessian herds and possible preventive measures. *Tierärztliche Umschau* 50, 250-255.
- Gu, B., Zhu, Y., Zhu, W., Miao, J., Deng, Y.E., Zou, S., 2009. Retinoid protects rats against neutrophil-induced oxidative stress in acute experimental mastitis. *International Immunopharmacology* 9, 223–229.
- Halliwell, B., 1991. Reactive oxygen species in living systems, source, biochemistry and role in human disease. *American Journal of Medicine* 91, 1422.
- Halliwell, B., 1994. Free radicals, antioxidants, and human disease, Curiosity, cause or consequence. *Lancet* 344, 721-724.
- Hicks, C.L., Korycka-Dahl, M., Richardson, T., 1975. Superoxide dismutase in bovine milk. *Journal of Dairy Science* 58, 796.
- Hogan, J.S., Smith, K.L., Weiss, W.P., Todhunter, D.A., Schockey, W.L., 1990 Relationships among vitamin E, selenium, and bovine blood neutrophils. *Journal of Dairy Science* 73, 2372–2378.
- Hogan, J.S., Weiss, W.P., Smith, K.L., 1993. Role of vitamin E and selenium in host defense against mastitis. *Journal of Dairy Science* 76, 2795–2803.
- Hogan, J.S., Weiss, W.P., Smith, K.L., 1995. Effects of *Escherichia coli* J5 vaccine on mild clinical coliform mastitis. *Journal of Dairy Science* 78, 285-290.
- Holbrook, J., Hicks, C.L., 1978. Variation of Superoxide Dismutase in Bovine Milk. *Journal of Dairy Science* 61, 1072-1077.
- Huie, R.E., Padmaja, S., 1993 The reaction of NO with superoxide. *Free Radical Research* 18, 195–199.
- Huijps, K., Hogeveen, H., 2008. Costs of mastitis, Facts and perception. *Journal of Dairy Research* 75, 113-120.
- Jukola, E., Hakkarainen, J., Saloniemi, H., 1996. Blood selenium, vitamin E, vitamin A, and  $\beta$ -carotene concentrations and udder health, fertility treatments and fertility. *Journal of Dairy Science* 79, 838-845.
- Jungi, T.W., 2000. Research from the Division of Immunology, production of nitric oxide, NO, by macrophages in ruminants. *Schweizer Archiv für Tierheilkunde* 142, 215–217.
- Kincaid, R.L., Hodgson, A.S., Riley, R.E., 1984. Supplementation of diets for lactating with zinc oxide and zinc methionine *Journal of Dairy Science Suppl.* 1,103.
- Kleczkowski, M., Klucinski, W., Shaktur, A., Sikora, J., 2005. Concentration of ascorbic acid in the blood of cows with subclinical mastitis. *Polish Journal of Veterinary Sciences* 8, 121–125.
- Knaapen, A.M., Seiler, F., Schilderman, P.A.E.L., Nehls, P., Bruch, J., Schins, R.P.F., Borm, P.J.A., 1999. Neutrophils cause oxidative DNA damage in alveolar epithelial cells. *Free Radical Biology and Medicine* 27, 234–240.
- Köhrle, J., Gärtner, R., 2009. Selenium and thyroid. *Best Practice and Research Clinical Endocrinology and Metabolism* 23, 815–27.
- Komine, K., Kuroishi, T., Komine, Y., Watanabe, K., Kobayashi, J., Yamaguchi, T., Kamata, S., Kumagai, K., 2004. Induction of nitric oxide production mediated by tumor necrosis factor alpha on staphylococcal enterotoxin c-stimulated bovine mammary gland cells. *Clinical and Diagnostic Laboratory Immunology* 11, 203–210.
- Lacasse, P., Lucy-Hulbert, J., Blais, S., 1997. Somatic cell production of the free radical nitric oxide during mastitis. *Livestock Production Science* 50, 168.
- Lindmark-Mansson, H., Akesson, B., 2000. Antioxidative factors in milk. *British Journal of Nutrition* 84, 103-110.
- MacMicking, J., Xie, Q., Nathan, C., 1997. Nitric oxide and macrophage function. *Annual Review of Immunology* 15, 323–350.
- McDowell, L.R., 1989. *Vitamins in Animal Nutrition*. Academic Press, Inc., San Diego, CA.
- Mee, J.F., O'Farrell, K., Rogers, P., 1993. Trace element deficiency in Irish dairy herds. *Irish Grassland and Animal Production Association Journal* 27, 29-40.
- Miller, J.K., Brzezinska-Slebodzinska, E., Madsen, F.C., 1993. Oxidative stress antioxidants and animal function. *Journal of Dairy Science* 76, 2812–2823.
- Miranda, M., Muriach, M., Almansa, I., Jareño, E., Bosch-Morell, F., Romero, F.J., Silvestre D., 2004. Oxidative status of human milk and its variations during cold storage. *BioFactors* 20, 129-137.
- Moilanen, E., Vapaatalo, H., 1995. Nitric oxide in inflammation and immune response. *Annals of Medicine* 27, 359–367.
- Mukherjee, R., 2008. Selenium and vitamin E increases polymorphonuclear cell phagocytosis and antioxidant levels during acute mastitis in riverine buffaloes. *Veterinary Research Communications* 32, 305–313.
- Naresh, R., Dwivedi, S.K., Swarup, D., Patra, R.C., 2002. Evaluation of ascorbic acid treatment in clinical and subclinical mastitis of Indian dairy cows. *Asian-Australasian Journal of Animal Science* 15, 905–911.
- Notebaert, S., Demon, D., Vanden Berghe, T., Vandenabeele, P., Meyer, E., 2008. Inflammatory mediators in *Escherichia coli*-induced mastitis in mice. *Comparative Immunology, Microbiology and Infectious Diseases* 31, 551–565.
- O'Flaherty, C., Beorlegui, N., Beconi, M.T., 2003. Participation of superoxide anion in the capacitation of cryopreserved bovine sperm. *International Journal of Andrology* 26, 109–114.
- Oldham E.R., Eberhart R.J., Muller L.D., 1991. Effects of supplemental vitamin A and  $\beta$ -carotene during the dry period and early lactation on udder health. *Journal of Dairy Science* 74, 3775-3781.
- Papp, L.V., Lu, J., Holmgren, A., Khanna, K.K., 2007. From selenium to selenoproteins, Synthesis, identity, and their role in human health. *Antioxid. Redox Signal.* 9,775–806.
- Pár, A., Jávör, T., 1984. Alternatives in hepatoprotection, cytoprotection- influences on mono-oxidase system-free

- radical scavengers a review. *Acta Physiologica Hungarica* 64, 409–423.
- Parantainen, J., Tenhunen, E., Kangasniemi, R., Sankari, S., Atroshi, F., 1987. Milk and blood levels of silicon, selenium status in bovine mastitis. *Veterinary Research Communications* 115, 467–477.
- Pavalata, L., Illek, J., Pechova, A., 2002. Selenium status of cattle in the Czech Republic. *Acta Veterinaria Brno* 71, 3-8.
- Popovic, Z., 2004. Performance and udder health status of dairy cows influenced by organically bound zinc and chromium. Ph.D. thesis, University of Belgrade.
- Przybylska, J., Albera, E., Kankofer, M., 2007. Antioxidants in Bovine Colostrum. *Reproduction in Domestic Animals* 42, 402-409.
- Ranjan, R., Swarup, D., Naresh, R., Patra, R.C., 2005. Enhanced erythrocytic lipid peroxides and reduced plasma ascorbic acid, and alteration in blood trace elements level in dairy cows with mastitis. *Veterinary Research Communications* 29, 27–34.
- Rice, D.A., Kennedy, S., 1988. Assessment of vitamin E, selenium and polyunsaturated fatty acid interactions in the aetiology of disease in the bovine. *Proceedings of the Nutrition Society* 47, 177-184.
- Riollet, C., Rainard, P., Poutrel, B., 2000. Differential induction of complement fragment C5a and inflammatory cytokines during intramammary infections with *Escherichia coli* and *Staphylococcus aureus*. *Clinical and Diagnostic Laboratory Immunology* 7, 161–167.
- Salman, S., Khol-Parisini, A., Schafft, H., Lahrssen-Wiederholt M., Hulan H.W., Dinse D., Zentek J., 2009. The role of dietary selenium in bovine mammary gland health and immune function. *Animal Health Research Review* 10, 21–34.
- Sauberlich, H.E., 1994. Pharmacology of vitamin C. *Annual Reviews in Nutrition* 14, 371–391.
- Scaletti, R.W., Hamilton, C.H., Harmon, R.J., 2002. Effect of copper source on resistance to coliform mastitis. *Journal of Dairy Science* 85, 375.
- Scaletti, R.W., Trammell, D.S., Smith, B.A., Harmon, R.J., 2003. Role of dietary copper in enhancing resistance to *Escherichia coli* Mastitis. *Journal of Dairy Science* 86, 1240–1249.
- Smith J.A., 1994. Neutrophils, host defense, and inflammation, a double-edged sword. *Journal of Leukocyte Biology* 56, 672–686.
- Smith K.L., Conrad H.R., Amiet B.A., 1985. Incidence of environmental mastitis as influenced by vitamin E and selenium. *Kiel. Milchwir tsch. Forschungsber* 37, 482.
- Smith, K.L., Harrison, J.H., Hancock, D.D., Todhunter, D.A., Conrad, H.R., 1984. Effect of vitamin E and selenium supplementation on incidence of clinical mastitis and duration of clinical symptoms. *Journal of Dairy Science* 67, 1293–1300.
- Smith, K.L., Hogan, J.S., Weiss, W.P., 1997. Dietary vitamin E and selenium affect mastitis and milk quality. *Journal of Animal Science* 75, 1659–1665.
- Sordillo, L.M., O'Boyle, N., Gandy, J.C., Corl, C.M., Hamilton, E., 2007. Shifts in thioredoxin reductase activity and oxidant status in mononuclear cells obtained from transition dairy cattle. *Journal of Dairy Science* 90, 1186–1192.
- Su, W.J., Chang, C.J., Peh, H.C., Lee, S.L., Huang, M.C., Zhao, X., 2002. Apoptosis and oxidative stress of infiltrated neutrophils obtained from mammary glands of goats during various stages of lactation. *American Journal of Veterinary Research* 63, 241-246.
- Suriyasathaporn, W., Vinitketkumnuen, U., Chewonarin, T., Boonyayatra, S., Kreausukon, K., Schukken, Y.H., 2006. Higher somatic cell counts resulted in higher malondialdehyde concentrations in raw cows' milk. *International Dairy Journal* 16, 1088-1091.
- Trigona, W.L., Mullarky, I.K., Cao, Y., Sordillo, L.M., 2006. Thioredoxin reductase regulates the induction of haem oxygenase-1 expression in aortic endothelial cells. *Biochemistry Journal* 394, 207–216.
- Wang, P.G., Xian, M., Tang, X., Wu, X., Wen, Z., Cai, T., Janczuk, A.J., 2002. Nitric oxide donors, chemical activities and biological applications. *Chemical Reviews* 102, 1091–1134.
- Weiss, W.P., Hogan, J.S., Smith, K.L., 1990. Relationships among selenium, vitamin E, and mammary gland health in commercial dairy herds. *Journal of Dairy Science* 73, 381.
- Weiss, W.P., Hogan, J.S., Smith, K.L., 2004. Changes in vitamin C concentrations in plasma and milk from dairy cows after an intramammary infusion of *Escherichia coli*. *Journal of Dairy Science* 87, 32–37.
- Weiss, W.P., Hogan, J.S., Todhunter, D.A., Smith, K.L., 1997. Effect of vitamin E supplementation in diets with a low concentration of selenium on mammary gland health of dairy cows. *Journal of Dairy Science* 80, 1728–1737.
- Zust, J., Hrovatin, B., Simundic, B., 1996. Assessment of selenium and vitamin E deficiencies in dairy herds and clinical disease in calves. *Veterinary Record* 139, 391-394.