

## Estimation of Total Saponins and Evaluate Their Effect on in vitro Methanogenesis and Rumen Fermentation Pattern in Wheat Straw Based Diet

Navneet Goel<sup>1</sup>, Sunil Kumar Sirohi<sup>1\*</sup>, Jaya Dwivedi<sup>2</sup>

<sup>1</sup>Nutrition Biotechnology Lab, Dairy Cattle Nutrition Division, National Dairy Research Institute, Karnal, 132001-Haryana, INDIA.

<sup>2</sup>Department of Chemistry, Banasthali University, Banasthali, 304022-Rajasthan, India.

(Received 26 November 2011/ Accepted 6 April 2012)

### Abstract

The present experiment was carried out to estimate the total saponins and evaluate their effect on methanogenesis and rumen fermentation by in vitro gas production techniques. Three plant material, rough chaff tree seed (*Achyranthus aspara*, T<sub>1</sub>), gokhru seed (*Tribulus terrestris*, T<sub>2</sub>) and Siris seed (*Albizia lebeck*, T<sub>3</sub>) were selected for present study. The total saponins content in T<sub>1</sub>, T<sub>2</sub> and T<sub>3</sub> were 45.75, 25.65 and 48.26% (w/w), respectively. Three levels of each saponins (3, 6 and 9% on DM basis) and wheat straw based (50R:50C) medium fiber diet (200±10 mg) were used for the evaluation of their effect on methanogenesis and rumen fermentation pattern. Results showed the maximum methane reduction (49.66% in term of mM/gDDM) and acetate propionate ration (35.08%) were found in T<sub>1</sub> at 6 and 3% levels. Result show that propionate production (mM/ml) was increased; protozoa population decreased (75%) significantly on addition with T<sub>3</sub> at 6% level. No significant variation was found in dry matter digestibility in all cases. The present results demonstrate that total saponins extracted from different herbal plants are a promising rumen modifying agent. They have the potential to modulate the methane production, dry matter digestibility and microbial biomass synthesis.

**Keywords:** Saponins; In vitro gas production technique; methane; IVDMD; rumen fermentation

### Introduction

The recent goal of ruminant microbiologists and nutritionists is to manipulate the ruminal microbial ecosystem to improve the feed conversion efficiency and reduce methane production. Most of the recent studies deal with the diversity analysis of rumen microbes especially methanogens (Chaudhary 2009; Chaudhary *et al.*, 2009, 2012) but the studies that deal with the manipulation in the rumen microbial ecosystem for increasing digestion and improving other nutritional parameters are still limited. Increasing fiber digestion, propionate production, yield and efficiency of microbial protein synthesis and decreasing methanogenesis, extensive dietary protein degradation and bacterial recycling by protozoa in rumen are some of the well recognized ways to improve efficiency of nutrient utilization and ruminant productivity under roughage based feeding systems. Recent studies

have shown that PSM at lower concentrations could be used to manipulate rumen fermentation and to improve animal productivity. Plant secondary metabolites present in different plant extract positively affected the ruminal methanogenesis (Sirohi *et al.*, 2009; Goel *et al.*, 2011). Although effort have also been made to select the potential inhibitors from chemical compounds available to check methanogenesis by computational means to get better results (Sharma *et al.*, 2011). Saponins, tannins and EO (Essential oil) are the group of secondary metabolites that have been widely investigated for their pro nutritional effects in ruminants. Among the all PSM, saponins attract the researchers due to their presence of large group of plants, their antimethanogenic activity and many beneficial effects on animal systems (Francis *et al.*, 2002a). Saponins are surface-active glycosides with detergent, wetting, emulsifying, and foaming properties (Wang *et al.*, 2005, Sarnthein *et al.*, 2004; Mitra *et al.*, 1997) and occurring widely in plants and some lower marine animals such as star fish and sea cucumber and bacteria (Riguera 1997; Yoshiki *et al.*, 1998). They are abundant in many

\*Corresponding author: Sunil Kumar Sirohi

Address: Nutrition Biotechnology Lab, Dairy Cattle Nutrition Division, National Dairy Research Institute, Karnal, 132001-Haryana, INDIA.

E-mail address: sirohisk@gmail.com

foods consumed by animals and man (Cheeke 1971). Methanolic extract of soapnut decreased *in vitro* methane production and increased molar proportion of propionate (Agarwal *et al.*, 2004). Saponin rich aqueous extract of *Asparagus adscendens* (satavar) root and aqueous methanol of *Moringa oleifera* (drumstick) seed decreased protozoal number and methanogenesis and increased yield and efficiency of microbial protein synthesis (Alexander, 2005).

## Materials and methods

### Procedure of saponins extracts preparation

Three plant parts, rough chaff tree seed (*Achyranthus aspara*, T<sub>1</sub>), gokhru seed (*Tribulus terrestris*, T<sub>2</sub>) and Siris seed (*Albizia lebbbeck*, T<sub>3</sub>) were used for the preparation of crude saponins extracts. Seeds of T<sub>1</sub> and T<sub>3</sub> were collected from the campus of National Dairy Research Institute, Karnal while T<sub>2</sub> seeds were purchased from local market of Karnal. Seeds of these plants were dried at 70°C and ground in mills to pass a 1 mm sieve. All plant parts were defatted with petroleum ether in a soxhlet apparatus.

Take 25 g fat free sample in 500 ml conical flask and add 250 ml absolute methanol (99.9%) in ratio of dry weight of the sample to methanol as 1:10. Flask was tightly sealed and kept in a shaker at 25°C and 120 rpm for 24 hour, followed by centrifuging the contents at 3500 rpm for 20 min. After centrifugation, methanol extract was filtered using Whatman filter paper No.1. The resulting methanolic extracts were evaporated to dryness under *in vacuo* condition using a rota-evaporator. After evaporation dried plant extract dissolve in minimum amount of distilled water (10 ml), transferred in to a separating funnel and extracted with equal volume of n-butanol (3 times). Again, solvent, n-butanol was evaporated under *in vacuo* condition not higher than 45°C. Dissolved the dried saponins content in 5-10 ml of distilled water and transferred the solution in to a separate pre-weighed container and freeze dried.

### Estimation of total saponins

Total saponins contents in different plant materials were estimated by prescribed colorimetric methods (Hiai *et al.*, 1976). Ten milligram of saponins ex-

tract was dissolve in 5 ml 80% aqueous methanol and 50 µl of solution was taken in different test tubes to which 0.25 ml of vanillin reagent (8%, w/v in 99.9% ethanol) was added. Test tubes were placed in ice-cold water bath and 2.5 ml of 72% (v/v) sulphuric acid was added slowly on the inner side of the wall. After mixing the content in each tube, these were left as such for 3 min. warmed the tubes to 60°C for 10 min using a water bath and then cooled them in ice-cold water bath. Absorbance was measured at 544 nm using spectrophotometer against the reagent blank and prepared the standards curve. Quillaja saponin (Sigma-Aldrich) was used as a reference standard (Shiau *et al.*, 2009) and the content of total saponins was expressed as Quillaja saponin equivalents (QS µg/mg extract).

### Preparation of diet

To evaluate the effect of different saponins extract, diet was prepared by taking roughage concentrate ratio of 50:50. The roughage part composed of wheat straw and the concentrate part composed of maize (33%), GNC (21%), mustard cake (12%), wheat bran (20%), deoiled rice bran (11%), mineral mixture (2%) and salt (1%) respectively.

### Treatments and experimental design

For the evaluation, stock solution (2mg/ml) of extracted saponins was prepared in distilled water and its different volumes were added in calibrated 100ml glass syringe containing 200±10 mg of milled (1mm) wheat straw based diet to have final concentrations of 3, 6 and 9% on DM basis. All the treatments were arranged in RBD with three replicates. Sets was also incubated devoid of substrate with and with out plant powder combinations which served as blanks for particular treatment and values were corrected for different parameters with these blanks.

### *In vitro* gas production

The incubation medium was prepared by prescribed method (Menke and Steingass, 1988). Rumen liquor was collected from a fistulated male buffalo (*Bubalus bubalis*) maintained on a standard diet (60 parts roughage: 40 parts concentrate) before morning feeding into an insulated flask and

brought into the laboratory. The rumen liquor filtered through four layers of muslin cloth and then the required amount of filtered rumen liquor used as a source of inoculum. Plungers of syringes applied with petroleum jelly for smooth movement and stop any leakage. The 30 ml incubation medium was dispensed anaerobically in each syringe, closed the clamps and incubated at  $39 \pm 0.5^\circ\text{C}$  for 24 h.

#### *Total gas production and methane estimation*

After 24 h incubation, total gas production was calculated by subtracting gas produced in blank syringe (containing no substrate, but only the inoculum and buffer) from total gas produced in the syringe containing substrate and inoculum and buffer. Methane content in fermentation gas was determined by gas chromatography (GC) using Nucon-5765 gas chromatograph equipped with flame ionization detector (FID) and stainless steel column packed with Porapak-Q (length 6'; o.d. 1/8" i.d. 2 mm; mesh range 80-100). Temperatures were 40, 50 and  $50^\circ\text{C}$ , in injector oven, column oven and detector, respectively and the flow rates of carrier gas (nitrogen), hydrogen and air were 30, 30 and 300 ml/min, respectively. For methane estimation, each gas sample (250  $\mu\text{l}$ ) was manually injected using Hamilton airtight syringe. Methane content in sample was calculated by external calibration, using a certified gases mixture with 50%  $\text{CH}_4$  and 50%  $\text{CO}_2$  (Spantech calibration gas, Surrey, England). The volume of methane produced was calculated as follows:

Methane production (ml) = Total gas produced (ml)  $\times$  % methane in the sample

#### *Partitioning factor and Microbial biomass yield*

The partitioning factor is calculated as the ratio of substrate truly degraded in vitro (mg) to the volume of gas (ml) produced by it. The microbial biomass (MBM) yield was calculated by using the degradability of substrate and gas volume and stoichiometrical factor (Blummel *et al.*, 1997).

$\text{MBM (mg)} = \text{Substrate truly degraded} - (\text{gas volume} \times \text{stoichiometrical factor})$

Where the stoichiometrical factor used was 2.25.

#### *Total volatile fatty acid (TVFA) estimation*

TVFA concentration (mM/100ml) in the supernatant was estimated according to prescribed

method (Barnet and Reid, 1957).

#### *Individual volatile fatty acid (IVFA) estimation*

Individual volatile fatty acid estimated by gas chromatograph according to the prescribed method (Erwin *et al.*, 1961).

#### *Estimation of ammonia nitrogen*

The supernatant of each syringe including that of blank was used for  $\text{NH}_3\text{-N}$  estimation. Supernatant (5 ml) was mixed with 1 N NaOH (2 ml) and steam passed on this using KEL PLUS - N analyzer (Pelican, India) and the  $\text{NH}_3$  evolved was collected in boric acid solution having mixed indicator and titrated against N/100  $\text{H}_2\text{SO}_4$ .

#### *Protozoa counting*

The protozoa in fermentation fluid were counted by Haemocytometer as per the prescribed method (Dehority, 1984).

#### *In vitro true DM degradability*

In vitro true DM degradability of feed sample of each syringe containing residues after incubation was estimated as per the prescribed method (Van soest *et al.*, 1991).

#### *Proximate analysis and Cell wall constituents*

The proximate analysis (Organic matter, Crude protein, Ether extract, Total Ash) of substrate was carried out as per the methods of AOAC (1995). The Neutral detergent fibre of substrates were determined according to prescribed method (Van Soest *et al.*, 1991) and other cell wall components like Acid detergent fiber (ADF) and Hemicellulose (HC) as per the method (Van Soest *et al.*, 1991).

#### **Statistical analysis**

Experimental data of different parameters were analyzed in randomized block design with three replicates for analysis of variance (Snedecor and Cochran, 1968).

#### **Results**

The physical and chemical composition of wheat

straws based medium fiber diet was presented in table 1. The OM, CP and NDF content of wheat straws based medium fiber diet were 878.4, 125.3 and 604.5 g/kg, respectively.

Table 1. Physical and chemical composition of wheat straw based diet used as substrate in in vitro incubation

Ingredient of diet							
Particulars	g/kg on DM basis						
Wheat straw	500						
Concentrate	500						
Ingredient of concentrate							
Particulars	g/kg on DM basis						
Maize	330						
Ground nut cake	210						
Mustard cake	120						
Wheat bran	200						
Deoiled rice bran	110						
Mineral mixture	20						
Salt	10						
Chemical constituents of diet (g/kg on DM basis)							
Diet	OM	CP	EE	NDF	ADF	HC	TA
R:C (50:50)	878.4	125.3	30.4	604.5	329.5	275.0	121.6

OM= Organic matter, CP= Crude protein, EE= Ether extract, NDF= Neutral detergent fiber, ADF= Acid detergent fiber, HC= Hemicelluloses, TA = Total ash

The total saponins content in *Achyranthus aspara* (T<sub>1</sub>), *Tribulus terrestris* (T<sub>2</sub>) and *Albizia lebbbeck* (T<sub>3</sub>) seeds were shown in table 2. T<sub>3</sub> contain highest total saponins content 48.26% (w/w) while T<sub>1</sub> and T<sub>2</sub> seeds have 45.75, 25.65% (w/w), respectively.

Effects of different doses of total saponins supplementation on in vitro rumen fermentation pattern and methane production of wheat straws based

Table 2. Total saponin contents in herbal plants parts (% w/w) as Quillaja saponin equivalents

Herbal Plants	Total saponins % (w/w)
<i>Achyranthus aspara</i> , (T <sub>1</sub> )	45.70
<i>Tribulus terrestris</i> , (T <sub>2</sub> )	25.65
<i>Albizia lebbbeck</i> , (T <sub>3</sub> )	48.26

medium fiber diet were shown in table 3 and table 4, respectively.

Methane production was reduced in each treatment and every level of doses. The highest inhibition of CH<sub>4</sub> (mM/gm digestible dry matter) formation occurred with T<sub>1</sub> (49.66%) at 6% level. A good reduction percentage in methane production were also observed in T<sub>2</sub> (47.61%) and T<sub>3</sub> (48.99%) at 9 and 6% levels, respectively. Significant reduction in protozoa population observed except T<sub>2</sub> treatment. Maximum 75% reduction of protozoa population was found in T<sub>3</sub> at 6% level. Among the all treatment and doses, total volatile fatty acids concentration (mM/100ml) was maximum increased (9.55%) in T<sub>1</sub> at 3% level. T<sub>3</sub> at 3% level showed the maximum increase in Propionate concentration (13.39%), it was also show the maximum reduction in acetate (26.56%) and butyrate (30.35%) concentration (mM/100ml). Reduction in acetate propionate ratio was highest in T<sub>1</sub> (35.08%) at 3% level. At 6% level of T<sub>2</sub> and T<sub>3</sub> were also show the good reduction percentage in acetate propionate ratio i.e. 24 and 21.23% respectively. Supplementation of total saponins at differ-

Table 3. Supplementation effect of different saponins extracts on in vitro rumen fermentation pattern and methane inhibition on wheat straw based diet (50R:50C)

Treatment	Dose (%DM Basis)	IVDMD%	CH <sub>4</sub> (ml/gm DM)	CH <sub>4</sub> (mM/gm DM)	MBM (mg)	PF	Protozoa (x10 <sup>4</sup> /ml)
Control	00	64.65±0.70	37.57±1.52	1.47±0.06	40.06±1.70	3.26±0.05	2.00±0.29
T <sub>1</sub>	3	70.80±0.64	24.14±0.29	0.94±0.01	50.47±0.72	3.50±0.04	1.67±0.17
T <sub>1</sub>	6	68.68±0.55	18.96±0.43	0.74±0.02	58.62±1.81	3.93±0.07	1.33±0.33
T <sub>1</sub>	9	67.09±1.24	24.47±0.73	0.96±0.03	50.93±0.22	3.63±0.03	1.00±0.29
T <sub>2</sub>	3	66.52±1.46	20.83±0.97	0.81±0.04	55.78±5.62	3.89±0.23	1.50±0.29
T <sub>2</sub>	6	66.13±0.85	21.40±0.91	0.84±0.03	52.38±3.02	3.74±0.14	1.25±1.14
T <sub>2</sub>	9	64.60±1.69	19.74±1.26	0.77±0.05	47.82±4.36	3.58±0.14	1.50±0.29
T <sub>3</sub>	3	64.10±1.11	21.82±0.37	0.85±0.02	45.69±1.88	3.50±0.05	0.67±0.17
T <sub>3</sub>	6	67.54±1.86	19.23±0.86	0.75±0.02	61.20±3.25	4.12±0.11	0.50±0.29
T <sub>3</sub>	9	72.49±3.80	22.44±1.01	0.88±0.04	61.36±11.17	3.93±0.39	1.00±0.29

T<sub>1</sub> = *Achyranthus aspara*, T<sub>2</sub> = *Tribulus terrestris*, T<sub>3</sub> = *Albizia lebbbeck*, IVDMD = In vitro dry matter digestibility, CH<sub>4</sub> = Methane, MBM = microbial biomass, PF = Partition factor

Table 4. Supplementation effect of different saponins extracts on in vitro IVFA and Acetate: propionate ratio on wheat straw based diet (50R:50C)

Treatment	Dose (% DM Basis)	Acetate (mM/100ml)	Propionate (mM/100ml)	Butyrate (mM/100ml)	A:P	TVFA (mM/100 ml)	NH <sub>3</sub> -N (mg/100ml)
Control	00	10.46±0.41	3.21±0.03	1.68±0.02	3.25±0.09	7.85±0.08	21.75±0.10
T <sub>1</sub>	3	7.68±0.07	3.64±0.06	1.17±0.04	2.11±0.07	8.60±0.10	18.20±0.16
T <sub>1</sub>	6	7.88±0.48	3.62±0.14	1.14±0.07	2.18±0.04	8.07±0.09	18.39±0.09
T <sub>1</sub>	9	7.87±0.25	3.56±0.12	1.19±0.06	2.20±0.07	7.95±0.10	18.48±0.16
T <sub>2</sub>	3	8.43±0.17	3.15±0.13	1.18±0.01	2.67±0.11	7.72±0.02	18.85±0.41
T <sub>2</sub>	6	8.57±0.09	3.47±0.03	1.37±0.01	2.47±0.07	8.13±0.06	18.67±0.25
T <sub>2</sub>	9	8.78±0.35	3.04±0.08	1.38±0.08	2.89±0.06	7.72±0.06	19.60±0.16
T <sub>3</sub>	3	10.14±0.14	3.37±0.06	1.50±0.03	3.01±0.08	8.08±0.02	18.01±0.25
T <sub>3</sub>	6	9.20±0.12	3.59±0.12	1.35±0.05	2.56±0.07	8.05±0.03	18.39±0.34
T <sub>3</sub>	9	9.29±0.25	3.22±0.08	1.33±0.11	2.88±0.02	7.93±0.11	17.55±0.28

T<sub>1</sub> = *Achyranthus aspara*, T<sub>2</sub> = *Tribulus terrestris*, T<sub>3</sub> = *Albizia lebbek*, A:P = Acetate to propionate ratio, TVFA = Total volatile fatty acid, NH<sub>3</sub>-N = Ammonia nitrogen

ent levels did not affect dry matter digestibility of wheat straw based medium fiber diet. The highest increase in dry matter digestibility was in T<sub>3</sub> (12%) at 9% level. The partition factor value were increased in all cases but the maximum raise was found in T<sub>3</sub> (26.38%) at 6% level. Among the all treatment, Microbial biomass (mg) was significantly increased in T<sub>3</sub> (53.17%) at 9% level and it was also increased more then 50% in case of T<sub>1</sub> at 6% level.

## Discussion

Ruminants produced methane by enteric fermentation, which is leads to a sever loss of feed energy. Methane is potent greenhouse gas; therefore, reducing methane production has significant economical and environmental benefits. Inclusion of T<sub>3</sub> at 6% level reduced the 24 h methane emission significantly. Inhibition of methanogenesis by T<sub>3</sub> was primarily due to their anti-protozoal activity. As some methanogens live in association with protozoa (New bold *et al.*, 1995; Tokura *et al.*, 1997), it was expected that reducing protozoa would also reduce methanogens, hence reducing methane production.

In present study, among the all treatment, T<sub>3</sub> showed the significantly reduced protozoa count (table 3). Saponins possibly bind with sterol of cell membrane of protozoa and change the permeability of cell membrane. Decreased protozoal counts with supplementation of saponins rich extract (Hristov *et al.*, 1999b; Kamra *et al.*, 2000) or saponin rich forages (Newbold *et al.*, 1997, Teferedegne *et al.*, 1999) or fruits (Thalib *et al.*, 1998; Hess *et al.*,

2003) have been reported. Increase in propionate (percent) and decrease in acetate (percent) and consequently decrease in acetate and propionate ratio by T<sub>3</sub> could be due to the high percentage of saponins and its inhibitory effect on protozoa, which is in agreement with previous studies (Wang *et al.*, 2000; Ye *et al.*, 2001; Lila *et al.*, 2003). Over all, the present results demonstrate that 6% T<sub>3</sub> is a promising level of supplementation than the other levels of T<sub>3</sub> and other saponins in 50:50 wheat straw based roughage concentrate diet. Total saponins of T<sub>3</sub> have the potential for use as a de-faunating agent and should be tested to find out a suitable dose to get maximum inhibition in methane emission without adversely affecting feed degradability.

## Acknowledgements

Authors would like to thank and acknowledge for the grant provided by NFBSRA, Ministry of Agriculture, ICAR, New Delhi-100012 to carry out this research work.

## References

- Agarwal, N., Srivastava, R.K., Kamra, D.N. 2004. Effects of soapnut extracts on in vitro methanogenesis, fermentation by rumen liquor of buffalo. In: Proceeding, New Dimensions of Animal Feeding to sustain Development and Competitiveness. V Biennial Conference of Animal Nutrition Association, 24-26 November, 2004, NIANP, Bangalore, India. pp 226.
- Alexander, G., 2005. Effect of plant extracts on rumen fermentation and nutrient utilization in sheep. Ph.D. Thesis, Indian Veterinary Research Institute (Deemed

- University), Izatnagar (UP).
- AOAC, 1995. Official Methods of Analysis. 16th ed. Association of Official Analytical Chemists, Arlington, VA.
- Arao, T., Udayama, M., Kinjo, J., Nohara, T., 1998. Preventive effects of Saponins from the *Pueraria lobata* root on in vitro immunological liver injury of rat primary hepatocyte cultures. *Planta Medica* 64, 413-416.
- Barnet, A.J.G., Reid, R.L., 1957. Studies on production of volatile fatty acids from grass by rumen liquor in artificial rumen IVFA production from fresh grass. *Journal of Agriculture Science* 48, 315.
- Blummel, M., Makkar, H.P.S., Becker, K., 1997. In vitro gas production: a technique revisited. *Journal of Animal Physiology and Animal Nutrition* 77, 24-34.
- Chao, A.C., Nguyen, J.V., Broughall, M., Recchia, J., Kensil, C.R., Daddona, P.E., Fix, J.A. 1998. Enhancement of intestinal model compound transport by DS-1, a modified *Quillaia* saponins. *Journal of Pharmaceutical Science* 87, 1395-1399.
- Chaudhary, P.P., Sirohi, S.K., Saxena, J., 2012. Diversity analysis of methanogens in rumen of *Bubalus bubalis* by 16S riboprinting and sequence analysis. *Gene* 493, 13-17.
- Chaudhary, P.P., Sirohi, S.K., 2009. Dominance of Methanobacterium phylotype in rumen (*Bubalus bubalis*) methanogens from India. *Letters in Applied Microbiology* 49, 274-277.
- Chaudhary, P.P., 2009. Methanobacterium phylotype are the dominant methanogen phylotype in the Murrah buffaloes. *Letters in Applied Microbiology* 48, 386.
- Cheeke P.R., 1971. Nutritional and physiological implications of saponins. A review: *Canadian Journal of Animal Science* 51, 621-632.
- Dehority, B.A., 1984. Evaluation of subsampling and fixation procedures used for counting rumen protozoa. *Applied and Environmental Microbiology* 48, 182-185.
- Erwin, E.S., Macro, G.A., Emery, E.M., 1961. Volatile fatty acid analysis of blood and rumen fluid by Gas chromatography. *Journal of Dairy Science* 44, 1768-1771.
- Francis G., Makkar H.P.S., Becker K., 2002a. Dietary supplementation with a *Quillaja* saponin mixture improves growth performance and metabolic efficiency in common carp (*Cyprinus carpio* L). *Aquaculture* 203, 311-320.
- Goel, N., Sirohi, S.K., Jaya Dwivedi., 2011. Studies on the effects of methanolic extract of *Cinnamomum zeylanicum* on in vitro methane inhibition and rumen fermentation pattern. *Journal of Chemical and Pharmaceutical Research* 3, 609-615.
- Hess, H.D., Kreuzer, M., Diaz, T.E., Lascano, C.E., Carulla, J.E., Solvia, C.R., 2003. Saponin rich tropical fruits affect fermentation and methanogenesis in faunated and defaunated fluid. *Animal Feed Science and Technology* 109, 79-94.
- Hess, H.D., Monsalve, L.M., Lascano, C.E., Carulla, J.E., Diaz, T.E., Kreuzer, M., 2003b. Supplementation of a tropical grass diet with forage legumes and *Sapindus saponaria* fruits: effects on in vitro ruminal nitrogen turnover and methanogenesis. *Australian Journal of Agriculture Research* 54, 703-713.
- Hiai, S., Oura, H., Nakajima, T., 1976. Color reaction of some saponin and saponins with vanillin and sulfuric acid. *Planta Medica* 29, 116-122.
- Hristov, A.N., McAllister, T.A., Van Herk, F.H., Cheng, K.J., Newbold, C.J., Cheeke, P.R., 1999b. Effect of *Yucca schidigera* on ruminal fermentation and nutrient digestion in heifers. *Journal of Animal Science* 77, 2554-2563.
- Jun, H.K., Park, K.Y., Jo, J.B., 1989. Inhibitory effects of Ginseng saponins on Aflatoxin production in culture. *Chemical Abstracts* 106, 116-199.
- Just, M. J., Reecsio, M.G., Gner, R.M., Cuellar, M.J., Marez, S., Bilia, A.R., Rios, J., 1998. Anti-inflammatory activity of unusual lupane saponins from *Buleurum fruticosum*. *Planta Medica* 64, 404-407.
- Kamra, D.N., Singh, R., Agarwal, N., Pathak, N.N., 2000. Soapnut (Reetha) as natural defaunating agent-its effect on rumen fermentation and in sacco degradability of jowar hay in buffaloes. *Buffalo Journal* 16, 99-104.
- Lila, Z.A., Mohammed, N., Kanda, S., Kamada, T., Itabashi, H., 2003. Effect of sarsaponin on rumen fermentation with particular reference to methane production in vitro. *Journal of Dairy Science* 86, 3330-3336.
- Lu, C.D., Tsai, L.S., Schaefer, D.M., Jorgensen, N.A., 1987. Alteration of fermentation in continuous culture of mixed rumen bacteria by isolated alfalfa saponins. *Journal of Dairy Science* 70, 799-805.
- Menke, K.H., Steingass, H., 1988. Estimation of the energetic feed value obtained by chemical analysis and in vitro gas production using rumen fluid. *Animal Research* 28, 7-55.
- Mitra, S., Dungan, S.R., 1997. Micellar properties of quillaja saponin. Effects of temperature, salt and pH on solution properties. *Journal of Agricultural and Food Chemistry* 45, 1587-1595.
- Newbold, C.J., Hassan, S.M.E., Wang, J., Ortega, M.E., Wallace, R.J., 1997. Influence of foliage from African multipurpose trees on activity of rumen protozoa and bacteria. *British Journal of Nutrition* 78, 237-249.
- Newbold, C.J., Lassalas, B., Jouany, J.P., 1995. The importance of methanogens associated with ciliate protozoa in ruminal methane production in vitro. *Letters in Applied Microbiology* 21, 230-234.
- Oakenfull, Fenwick, 1981. Saponin content of soybeans and some commercial soybean products. *Journal of the Science of Food and Agriculture* 32, 273-278.
- Okubo, K., Kudou, S., Uchida, T., Yoshiki, Y., Yoshikoshi, M., Tonomura, M., 1994. Soyabean saponin and Isoflavonoids: structure and antiviral activity against Human Immunodeficiency virus in vitro. *ACS Symp. Ser.* p.546.
- Patra, A.K., Kamra, D.N., Agarwal, N., 2004. Effect of leaf extracts on methanogenesis rumen fermentation and enzyme activities by in vitro gas production test. In: *New dimensions of animal feeding to sustain development and competitiveness. Proceeding V Biennial Conference of Animal Nutrition Association*, 24-26 November, 2004, NIANP, Bangalore, India. pp, 227.
- Price, Fenwick, 1990. The Chemistry and Biological significance of saponins in foods and feeding stuffs. *International Reviews in Food Science and Nutrition* 157-162.

- Riguera R., 1997. Isolating bioactive compounds from marine organisms. *Journal of Marine Biotechnology* 5, 187–193.
- Sarnthein-Graf, C., La Mesa, C., 2004. Association of saponins in water and water gelatine mixtures. *Thermochimica Acta* 418, 79–84.
- Sharma, A., Chaudhary, P.P., Sirohi, S.K., Saxena J., 2011. Structure modeling and inhibitor prediction of NADP oxidoreductase enzyme from *Methanobrevibacter smithii*. *Bioinformation* 6, 15-19.
- Shiau, I.L., Shih, T.L., Wang, Y.N., Chen, H.T., Lan, H.F., Lin, H.C., Yang, B.Y., Ko, C.H., Murase, Y., 2009. Quantification for saponin from a soapberry in cleaning products by a chromatographic and two colorimetric assays. *Journal of the Faculty of Agriculture Kyushu University* 54, 215–221.
- Sirohi, S.K., Pandey, N., Goel, N., Singh, B., Mohini, M., Pandey, P., Chaudhary P.P., 2009. Microbial Activity and Ruminal methanogenesis as affected by Plant Secondary Metabolites in Different Plant Extracts. *International Journal of Environmental Science and Engineering* 1, 52-58.
- Snedecor, G.W., W.G. Cochran., 1968. *Statistical Methods*, 5th ed. Iowa State Univ. Press, Ames., I.A.
- Teferedegne, B., McIntosh, F., Osuji, O.P., Odenyo, A., Wallace, R.J., Newbold, C.J., 1999. Influence of foliage from different accessions of the sub-tropical leguminous tree, *Sesbania sesban*, on ruminal protozoa in Ethiopian and Scottish sheep. *Animal Feed Science and Technology* 78, 11–20.
- Thalib, A., Widiawati, Y., Hamid, H., Suherman, D., Newbold, C.J., 1998. Influence of saponin/sapogenins on the bacteriolytic activity of ciliate protozoa from the rumen. *Proceeding British Society of Animal Science*, 122.
- Tokura, M., Ushida, K., Miyazaki, K., Kojima, Y., 1997. Methanogens associated with rumen ciliates. *FEMS Microbiology Ecology* 22, 137–143.
- Tschesche, R., Wulff, G., 1973. Chemistry and Biology of Saponins. *Fortschr Chem Org Naturst*, 30, 461.
- Van Soest, P.J., Robertson, J.B., Lewis, B.A., 1991. Methods for dietary fiber, neutral detergent fiber and non-starch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 74, 3583–3597.
- Wallace, R.J., Arthaud, L., Newbold, C.J., 1994. Influence of *Yucca schidigera* extract on ruminal ammonia concentrations and ruminal microorganism. *Applied and Environmental Microbiology* 60, 1762–1767.
- Wang, Y., Greer, D., McAllister, T.A., 2005. Effect of a saponin-based surfactant on water absorption, processing characteristics and in vitro ruminal fermentation of barley grain. *Animal Feed Science and Technology* 118, 255–266.
- Wang, Y., McAllister, T.A., Yanke, L.J., Cheeke, P.R., 2000. Effects of steroidal saponins from *Yucca schidigera* extract on ruminal microbes. *Journal of Applied Microbiology* 88, 887–896.
- Wang, Yu, Xi, McAllister, T.A., Yanke, L.J., Xu, Z.J., Cheeke, P.R., Cheng, K.J., Wang, Y.X., Cheng, K.J., 2000. In vitro effects of steroidal saponins from *Yucca schidigera* extract on rumen microbial protein synthesis and ruminal fermentation. *Journal of the Science of Food and Agriculture* 80, 2114–2122.
- Ye, Junan, Liu, JianXin, Shi, Z.Q., Jiang, L.S., Liu, J.X., 2001. Effect of tea saponins on fermentation of cultured rumen fluid. *Chinese Journal of Animal Science* 37, 29–30.
- Yoshiki Y., Kudou S., Okubo K., 1998. Relationship between chemical structures and biological activities of triterpenoid saponins from soybean (Review). *Bioscience Biotechnology and Biochemistry* 62, 2291–2299.
- Zhang, S., Hu, Z., 1985. Anti-ulcerogenic effects of Ginseng flower saponins in the rat *Zhongyao Tongbao* 10: 331 *Chemical Abstracts* 104,512.